

Appearance of vortices and monopoles in a decomposition of SU(2) Yang-Mills field

A. Mohamadnejad* and S. Deldar†

Department of Physics, University of Tehran,

P.O. Box 14395/547, Tehran 1439955961, Iran

Abstract

We propose a decomposition for SU(2) Yang-Mills field in the low energy limit. Motivated by Abelian dominance, we suppose that in the infrared regime of the SU(2) gauge theory, the field strength tensor can be obtained by multiplying two parts: $G_{\mu\nu}$ and $\mathbf{n}$, where $G_{\mu\nu}$ is a space-time tensor and the second part, $\mathbf{n}$, is an isotriplet unit vector field which gives the Abelian direction at each space-time point, $\mathbf{G}_{\mu\nu} = G_{\mu\nu} \mathbf{n}$. By considering some physical assumptions, this leads to a decomposition for the SU(2) Yang-Mills field. It seems that by this type of decomposition, both monopoles and vortices appear at the same time. In addition, in the presence of vortices, Dirac quantization condition is manifested with a rescaled electric charge, as well.

PACS numbers: 11.15.Ha, 12.38.Aw, 12.38.Lg, 12.39.Pn

*a.mohamadnejad@ut.ac.ir

†sdeldar@ut.ac.ir

I. INTRODUCTION

It is stated that ultraviolet and infrared limits of a Yang-Mills theory characterize different phases. The qualitative picture of this assertion is built especially by 't Hooft and Polyakov [1]. In the ultraviolet limit, the Yang-Mills theory is asymptotically free and perturbative methods are adequate. This limit describes the interaction between massless gluons which corresponds to the transverse polarizations of the gauge field A_μ . At Low energies the Yang-Mills theory becomes strongly coupled and perturbative techniques fail and nonperturbative methods must be developed. On the other hand, Yang-Mills theories in the low energy limit must exhibit color confinement. Therefore, describing the confinement problem needs nonperturbative methods and it has long been argued that the confinement can take place through the condensation of monopoles which leads to the dual Meissner effect in a dual superconductor [2]. For a review of the dual superconductor picture, see [3].

Non-perturbative effects in the low energy limit can be represented effectively by the topological structures of the gauge theory such as vortices and monopoles. The underlying gauge symmetry can be represented by the nontrivial topological degrees of freedom. There are some methods for extracting these topological degrees of freedom in the pure Yang-Mills theory. Examples are: Abelian projection [4] and field decomposition [5–7]. Abelian projection is a partial gauge fixing in which the projected gauge fields contain singularities interpreted as topological defects. In the second method, Abelian decomposition, one can do the same thing without gauge fixing [7]. The theories which result from these decompositions consist of a well-defined and self-consistent subset of a non-Abelian gauge theory for a given symmetry group. These theories are restricted and the dynamical degrees of freedom are reduced, providing us with a self-consistent but nontrivial subset of the original gauge theory. As a result of these methods, the original Yang-Mills theory turns into electrodynamics with magnetic monopoles, and they lead to Abelian dominance [8] and magnetic monopole dominance [9]. Indeed, in both Abelian projection [11] and field decomposition approach [12], one can obtain Wu-Yang magnetic monopoles [10], and based on their condensation the potential of a static quark-antiquark pair is derived. In agreement with Lattice results, this potential is composed of two parts: the first part is a Yukawa term that dominates the ultraviolet limit or the short range limit, and the second part is a linear term responsible for the confinement that dominates the low energy limit at large distances.

Even though the gauge field A_μ is a proper order parameter for describing the theory in its ultraviolet limit, in the infrared limit with monopole condensation some other parameters become more appropriate. Therefore, one can decompose the Yang-Mills field to new collective variables which are more appropriate for describing the long distance theory compared with the gauge field A_μ which corresponds to the short distance spectrum. Decomposing the Yang-Mills fields has been done by various methods. These decompositions pursue different purposes in particular in connection with the issue of quark confinement in quantum chromodynamics (QCD). In Cho method [5], the decomposition leads to a restricted theory with four dynamical degrees of freedom while an $SU(2)$ Yang-Mills field has six dynamical degrees of freedom. Hence, it seems that in the infrared limit, some degrees of freedom do not play considerable roles. Thus, the mechanism of color confinement in restricted theories will remain valid in the original one which is no longer restricted [5]. On the other hand, there are six degrees of freedom in the Faddeev-Niemi method [6]. One can obtain Skyrme-Faddeev Lagrangian by integrating out some of them [13]. In addition, Faddeev and Niemi have investigated the possibility that the low energy spectrum of pure Yang-Mills theory can be constructed of closed and knotted strings as stable solitons [6]. They derive an off-shell generalization of their decomposition [14] and they also present a new decomposition that realizes explicitly a symmetry between electric and magnetic variables, suggesting a duality picture between the corresponding phases [15].

In this paper, motivated by Abelian dominance for the infrared limit of the Yang-Mills theory, we propose a decomposition which has the same field strength tensor form as Cho theory, $\mathbf{G}_{\mu\nu} = G_{\mu\nu}\mathbf{n}$, but with three degrees of freedom. It seems that this decomposition describes the low energy limit including both vortices and monopoles as the topological structures that dominate the non-perturbative regime of the theory. We show that our decomposition is similar to the Faddeev-Niemi decomposition if we add a constraint to their decomposition. In addition, this decomposition manifests the Dirac quantization condition, but with a rescaled electric charge.

In the next section, we introduce our decomposition by defining $\mathbf{G}_{\mu\nu}$, $\mathbf{G}_{\mu\nu} = G_{\mu\nu}\mathbf{n}$, of the Cho decomposition. In this decomposition there exist three dynamical degrees of freedom. In sec. III, we discuss restricted $U(1)$ gauge theory to show how it leads to a string-like (vortex) object in a similar manner to Cho procedure for $SU(2)$ gauge theory supplying the Wu-Yang monopole. In sec. IV, we see how monopoles appear in the theory in addition to

vortices. Dirac quantization condition is obtained when the vortex configurations appear in the theory, but with a rescaled electric charge. Finally, the summary and the discussions are given in sec. V.

II. SU(2) YANG-MILLS FIELD IN THE LOW ENERGY LIMIT

In this paper, to avoid unnecessary complications, we concentrate on the the SU(2) gauge group which is the simplest non-Abelian group. Motivated by Cho decomposition and Abelian dominance, we assume that within a good approximation, the following form of the field strength tensor $\mathbf{G}_{\mu\nu}$, is dominant in the infrared limit of the SU(2) Yang-Mills theory,

$$\mathbf{G}_{\mu\nu} = G_{\mu\nu}\mathbf{n}. \quad (1)$$

$G_{\mu\nu}$ is a colorless tensor and $\mathbf{n}$ is a three components unit vector field pointing into the color direction. A natural way to make the decomposition, is construction of an orthogonal basis for the color space by $\mathbf{n}$ and its derivatives, for example: $\mathbf{n}$, $\partial_\mu\mathbf{n}$, and $\mathbf{n} \times \partial_\mu\mathbf{n}$ shown in Fig. (1):

$$\mathbf{n} \cdot \partial_\mu\mathbf{n} = \mathbf{n} \cdot (\mathbf{n} \times \partial_\mu\mathbf{n}) = \partial_\mu\mathbf{n} \cdot (\mathbf{n} \times \partial_\mu\mathbf{n}) = 0.$$

Then we expand the SU(2) Yang-Mills field as the following

$$\mathbf{A}_\mu = C_\mu\mathbf{n} + \phi_1\partial_\mu\mathbf{n} + \phi_2\mathbf{n} \times \partial_\mu\mathbf{n}, \quad (2)$$

where C_μ , ϕ_1 , and ϕ_2 are the coefficients of the expansion. Note that ϕ_1 and ϕ_2 are not independent of the direction μ . In general Eq. (2) above can be written as

$$\mathbf{A}_\mu = C_\mu\mathbf{n} + \phi_1^{(\mu)}\partial_\mu\mathbf{n} + \phi_2^{(\mu)}\mathbf{n} \times \partial_\mu\mathbf{n}, \quad (No\ sum\ over\ \mu)$$

Of course, it could be for some dynamical reason, that for Yang-Mills fields in the infrared regime $\phi_1^{(0)} = \phi_1^{(1)} = \phi_1^{(2)} = \phi_1^{(3)}$ and $\phi_2^{(0)} = \phi_2^{(1)} = \phi_2^{(2)} = \phi_2^{(3)}$. On the other hand, we know that an arbitrary SU(2) gauge field $\mathbf{A}_\mu$ has six dynamical degrees of freedom. Therefore, we suppose that ϕ_1 and ϕ_2 are independent of the direction μ so that totally, we have six dynamical degrees of freedom for $\mathbf{A}_\mu$, two for C_μ , two for $\mathbf{n}$, and two for ϕ_1 and ϕ_2 . This idea has been used by many authors including Faddeev and Niemi [6].




FIG. 1: Constructing an orthogonal basis for the SU(2) color space at each space-time point

Because of Eq. (1), C_μ , ϕ_1 , and ϕ_2 are not independent. In the following, we find the relation between them. For SU(2) field strength tensor, we have

$$\mathbf{G}_{\mu\nu} = \partial_\mu \mathbf{A}_\nu - \partial_\nu \mathbf{A}_\mu + g \mathbf{A}_\mu \times \mathbf{A}_\nu. \quad (3)$$

Substituting Eq. (2) in (3) and by considering the fact that we want the $G_{\mu\nu}$ has the form of Eq. (1), these relations between ϕ_1 , ϕ_2 , and C_μ are obtained:

$$\begin{aligned} \partial_\mu \phi_1 - C_\mu (1 + g\phi_2) &= 0, \\ \partial_\mu \phi_2 - gC_\mu \phi_1 &= 0. \end{aligned} \quad (4)$$

Changing the variables,

$$\begin{aligned} \phi_1 &= \frac{\rho}{g^2}, \\ 1 + g\phi_2 &= \frac{\sigma}{g}, \end{aligned} \quad (5)$$

where ρ and σ are real scalar fields and applying these new variables to Eq. (4), one obtains:

$$\begin{aligned} \partial_\mu \rho - gC_\mu \sigma &= 0, \\ \partial_\mu \sigma + gC_\mu \rho &= 0. \end{aligned} \quad (6)$$

The above equations can be written in a compact form:

$$D_\mu \phi = 0,$$

where D_μ is the covariant derivative, $D_\mu = \partial_\mu + igC_\mu$, and ϕ is a complex scalar field, $\phi = \rho + i\sigma$. Notice that the coupling between the scalar field ϕ , and the Abelian gauge

field C_μ , is the same as the coupling of the original SU(2) gauge theory, g . Therefore, we conclude that for satisfying Eq. (1) we must have the following decomposition for the SU(2) Yang-Mills field

$$\mathbf{A}_\mu = C_\mu \mathbf{n} + \frac{1}{g} \partial_\mu \mathbf{n} \times \mathbf{n} + \frac{\rho}{g^2} \partial_\mu \mathbf{n} + \frac{\sigma}{g^2} \mathbf{n} \times \partial_\mu \mathbf{n}, \quad (7)$$

with a constraint,

$$D_\mu \phi = (\partial_\mu + igC_\mu)(\rho + i\sigma) = 0. \quad (8)$$

This is our proposal for decomposing the SU(2) Yang-Mills Field. It looks like the Faddeev-Niemi decomposition [6], but with Eq. (8) as a constraint. This is the equation which shows the dependency of ρ , σ , and C_μ to each other.

A trivial solution for Eq. (8) is

$$\rho = \sigma = 0, \quad (9)$$

which leads to Cho decomposition with four dynamical degrees of freedom, two for $\mathbf{n}$ and two for C_μ corresponding to two polarization. The field strength tensor for Cho decomposition is:

$$\mathbf{G}_{\mu\nu} = \{(\partial_\mu C_\nu - \partial_\nu C_\mu) + (-\frac{1}{g} \mathbf{n} \cdot (\partial_\mu \mathbf{n} \times \partial_\nu \mathbf{n}))\} \mathbf{n}. \quad (10)$$

Equation (8) is familiar, indeed, Cho found his decomposition for SU(2) Yang-Mills field by a similar condition,

$$\nabla_\mu \mathbf{n} = (\partial_\mu + g \mathbf{A}_\mu \times) \mathbf{n} = 0 \Rightarrow \mathbf{A}_\mu = C_\mu \mathbf{n} + \frac{1}{g} \partial_\mu \mathbf{n} \times \mathbf{n}, \quad (11)$$

where $C_\mu = \mathbf{A}_\mu \cdot \mathbf{n}$. Hence, it seems that in the infrared limit of an Abelian or non-Abelian Yang-Mills theory, we have a Yang-Mills field so that it obliges the covariant derivative of scalar fields to be zero.

In the next section we find a non-trivial solution for Eq. (8) and we propose a decomposition for U(1) gauge field, C_μ , in the infrared regime.

III. RESTRICTED U(1) GAUGE THEORY AND VORTICES

In the previous section, we have proposed a decomposition, with a constraint. This constraint, Eq. (8), is worth studying independently. We show how this constraint restrict the Abelian U(1) gauge theory and leads to the appearance of string-like (vortex) objects. It

is similar to the case where the condition of Eq. (11) leads to the appearance of monopoles. Equation (8) implies,

$$\partial_\mu(\rho^2 + \sigma^2) = 0 \quad \Rightarrow \quad \phi^* \phi = \rho^2 + \sigma^2 = a^2, \quad (12)$$

where a is a constant. Equation (8) can be solved exactly for C_μ ,

$$C_\mu = \frac{1}{ga^2}(\sigma \partial_\mu \rho - \rho \partial_\mu \sigma). \quad (13)$$

The above equations show that the original gauge field, C_μ , which has two dynamical degrees of freedom has been restricted and it has one dynamical degree of freedom because it depends on ρ and σ which are not independent, shown in Eq. (12). Therefore, we have totally three dynamical degrees of freedom: two for $\mathbf{n}$ and one for C_μ or ϕ .

The field strength tensor can be written in terms of electric and magnetic field strength tensors, $F_{\mu\nu}$ and $H_{\mu\nu}$, respectively:

$$\mathbf{G}_{\mu\nu} = (F_{\mu\nu} + H_{\mu\nu})\mathbf{n}, \quad (14)$$

where

$$\begin{aligned} F_{\mu\nu} &= \partial_\mu C_\nu - \partial_\nu C_\mu, \\ H_{\mu\nu} &= -\frac{1}{g'}\mathbf{n} \cdot (\partial_\mu \mathbf{n} \times \partial_\nu \mathbf{n}). \end{aligned} \quad (15)$$

Notice that since $\rho^2 + \sigma^2 = a^2$, the contributions of ρ and σ are included in the new coupling g' , where

$$\frac{1}{g'} = \frac{1}{g} - \frac{a^2}{g^3}. \quad (16)$$

We take $a \leq g$ in order to have $g' \geq 0$. Note that the coupling constant increases in the low energy limit, $g' \geq g$, which is corresponding to the behavior of the coupling constant in Yang-Mills theories.

Now, we discuss the physical properties which can be obtained from $F_{\mu\nu}$ in Eq. (15). In the next section we study $H_{\mu\nu}$ of Eq. (15) and its properties.

Vortices which are the classical string-like objects, appear in this theory as topological singularities of the scalar field $\phi = (\rho, \sigma)$. Vortices are described by the homotopy class of a mapping $\Pi_1(S^1)$ of the spatial circle S_R^1 to the coset space $S = U(1)$ of the internal space. Now to define this mapping, one needs a two components scalar field in the theory, at least

on S_R^1 . The scalar field in our decomposition can be used to define the mapping $\Pi_1(S^1)$. We define the topological charge of the restricted U(1) theory by the homotopy class of the mapping $\Pi_1(S^1)$ given by (ρ, σ) :

$$(\rho, \sigma); \quad S_R^1 \rightarrow S^1 = U(1). \quad (17)$$

With this opening comment we show how to extract vortices. Notice that the homotopy class $\Pi_1(S^1)$ defined by "two-dimensional hedgehog ansatz" must describe the vortex with a unit flux tube:

$$(\rho, \sigma) = a \frac{\vec{r}}{r} = a(\cos(\varphi), \sin(\varphi)), \quad (18)$$

where φ is the azimuthal circular coordinate of S_R^1 . Using Eq. (18) in Eq. (13) one obtains

$$\begin{aligned} C_\mu &= -\frac{1}{g} \partial_\mu \varphi, \\ \implies C_r &= C_z = 0, \quad C_\varphi = -\frac{1}{gr}. \end{aligned} \quad (19)$$

The magnetic field $\vec{B}$ is obtained as the following

$$\begin{aligned} \vec{B} &= \vec{\nabla} \times \vec{C} = \hat{r} \left(\frac{1}{r} \frac{\partial C_z}{\partial \varphi} - \frac{\partial C_\varphi}{\partial z} \right) + \hat{\varphi} \left(\frac{\partial C_r}{\partial z} - \frac{\partial C_z}{\partial r} \right) \\ &\quad + \hat{k} \frac{1}{r} \left(\frac{\partial(r C_\varphi)}{\partial r} - \frac{\partial C_r}{\partial \varphi} \right) = 0. \end{aligned} \quad (20)$$

Indeed, in general we have

$$(\rho, \sigma) = a(\cos(\alpha), \sin(\alpha)) \Rightarrow C_\mu = -\frac{1}{g} \partial_\mu \alpha \Rightarrow F_{\mu\nu} = 0. \quad (21)$$

The above calculations is true for every place in space, but not on the z axis where $r = 0$. The magnetic flux passing through the closed curve C in Fig. (2), is not zero:

$$\begin{aligned} \phi_B &= \int_S \vec{B} \cdot \vec{ds} = \int_S (\vec{\nabla} \times \vec{C}) \cdot \vec{ds} = \oint_C \vec{C} \cdot \vec{dl} \\ &= \int_0^{2\pi} -\frac{1}{gr} r d\varphi = -\frac{2\pi}{g}. \end{aligned} \quad (22)$$

It shows that on the z axis the magnetic field is singular as well as C_φ in Eq. (19). So, although the magnetic field is zero everywhere, there exist an infinite magnetic field on the z axis, responsible for the magnetic flux of Eq. (22) which is an evidence of a string-like




FIG. 2: The magnetic field is zero everywhere but not on the z axis

object (vortex) lying on the z axis. One can obtain the magnetic field from Eq. (22)

$$\begin{aligned}
 \int_S \vec{B} \cdot d\vec{s} &= \int_0^R \int_0^{2\pi} B r d\theta dr = -\frac{2\pi}{g} \\
 \Rightarrow \int_0^R B r dr &= -\frac{1}{g} \\
 \Rightarrow B &= -2\frac{\delta(r)}{gr},
 \end{aligned} \tag{23}$$

therefore,

$$\vec{B} = B\hat{k}. \tag{24}$$

One can find out all the homotopically inequivalent classes of the mapping (17) and the corresponding vortex configurations by the following replacement

$$\varphi \rightarrow n\varphi. \tag{25}$$

Then the scalar field ϕ describes all homotopically inequivalent mapping of (17) with the homotopy class Z

$$Z = n \quad (n \text{ integer}), \tag{26}$$

corresponding to

$$C_r = C_z = 0, \quad C_\varphi = -\frac{n}{gr}, \tag{27}$$

for the vector potential, and

$$\vec{B} = -2\frac{n\delta(r)}{gr}\hat{k}, \tag{28}$$

for the magnetic field. The magnetic flux is:

$$\phi_B = -\frac{2\pi n}{g}. \quad (29)$$

IV. MONOPOLES AND RESCALED DIRAC QUANTIZATION CONDITION

In addition to the vortices which results from our decomposition, monopoles also emerge from $H_{\mu\nu}$ of Eq. (15). Up to now, we have proposed a decomposition (7)

$$\mathbf{A}_\mu = C_\mu \mathbf{n} + \frac{1}{g} \partial_\mu \mathbf{n} \times \mathbf{n} + \frac{\rho}{g^2} \partial_\mu \mathbf{n} + \frac{\sigma}{g^2} \mathbf{n} \times \partial_\mu \mathbf{n},$$

where the Abelian part C_μ , is also decomposed via Eq. (13), we call this decomposition a "double decomposition",

$$C_\mu = \frac{1}{ga^2} (\sigma \partial_\mu \rho - \rho \partial_\mu \sigma)$$

with the condition of Eq. (12):

$$\rho^2 + \sigma^2 = a^2.$$

We have obtained the field strength tensor,

$$\mathbf{G}_{\mu\nu} = (F_{\mu\nu} + H_{\mu\nu}) \mathbf{n},$$

where the first term, $F_{\mu\nu}$, leads to the appearance of the vortices. By studying the second term, we examine the monopole configurations which reproduce the Dirac quantization condition but with a rescaled electric charge, g' .

We know that for SU(2) gauge theory, the magnetic charge which is a topological charge, can be described by the homotopy class of the mapping $\Pi_2(S^2)$ of the two-dimensional sphere S_R^2 to the coset space $S^2 = SU(2)/U(1)$ of the internal space. To obtain the magnetic field, we choose a hedgehog configuration for $\mathbf{n}$:

$$\mathbf{n} = \frac{r^a}{r} = (\sin\alpha \cos\beta, \sin\alpha \sin\beta, \cos\alpha) \quad (30)$$

where

$$\alpha = \theta, \quad \beta = m\varphi.$$

θ and φ are the angular spherical coordinates of S_R^2 , and m is an integer number. The magnetic field can be obtained,

$$\begin{aligned} \vec{B} &= B \hat{r}, \\ B &= H_{\theta\varphi} = -\frac{1}{g'} \mathbf{n} \cdot (\partial_\theta \mathbf{n} \times \partial_\varphi \mathbf{n}) = -\frac{m}{g'} \frac{1}{r^2}, \end{aligned} \quad (31)$$

where

$$\partial_\theta = \frac{\partial}{r\partial\theta}, \partial_\varphi = \frac{\partial}{r\sin\theta\partial\varphi}.$$

From the Eq. (31):

$$\vec{\nabla} \cdot \vec{B} = -\frac{m}{g'} 4\pi \delta(\vec{r}), \quad (32)$$

which means that there is a magnetic monopole in the origin. Comparing Eq. (32) with the Gauss equation for a magnetic monopole charge, $-g_m$:

$$\vec{\nabla} \cdot \vec{B} = -4\pi g_m \delta(\vec{r}),$$

one gets

$$g'g_m = m. \quad (33)$$

This is the Dirac quantization condition, but with a rescaled electric charge g' . The original Dirac quantization condition, for Wu-Yang monopole or 't Hooft-Polyakov monopole, is

$$gg_m = m, \quad (34)$$

where g is the color electric charge and g_m is the magnetic color charge. But, for our decomposition, the relation between g and g_m is

$$gg_m = m(1 - (\frac{a}{g})^2). \quad (35)$$

If a goes to zero, the familiar Dirac quantization condition is restored. In addition, this new decomposition reduces to the Cho decomposition where the U(1) field is no longer decomposed, and consequently no vortices appear. So, it seems that the presence of vortices influence the Dirac quantization condition by rescaling the electric coupling or charge.

V. SUMMARY AND DISCUSSION

We have looked for a suitable decomposition of the SU(2) Yang-Mills field in the infrared region that would help to study the vacuum structure, in particular topological objects, like vortices and monopoles, believed in some models to be related to the phenomenon of color confinement. We conjecture that for the infrared limit of the SU(2) Yang-Mills theory, the field strength tensor which is a three-dimensional vector in the internal space can be obtained by multiplying two parts, $\mathbf{G}_{\mu\nu} = G_{\mu\nu}\mathbf{n}$. $G_{\mu\nu}$ is a tensor with the space-time indices

and the second part $\mathbf{n}$, is a three components unit vector field which selects the Abelian direction at each space-time point. This conjecture is motivated by Abelian dominance and the form of the field strength tensor of the Cho decomposition. We propose a decomposition corresponding to this conjecture. It looks like Faddeev-Niemi decomposition, but with the constraint $D_\mu\phi = (\partial_\mu + igC_\mu)\phi = 0$. This constraint is similar to the constraint of Cho decomposition $D_\mu\mathbf{n} = (\partial_\mu + g\mathbf{A}_\mu \times)\mathbf{n} = 0$. It seems that vanishing of the covariant derivative of scalar field in an Abelian or non-Abelian Yang-Mills theory has something to do with the low energy limit and the vacuum structure of the theory. As a result of this constraint in Abelian case, string-singularities or vortices can appear in the theory. $F_{\mu\nu}$ is zero everywhere except on the location of strings. However, for SU(2) gauge theory which is a non-Abelian theory the constraint leads to point-singularities (monopoles) and non-zero $G_{\mu\nu}$. Hence, the vacuum structure of a non-Abelian theory is richer than the Abelian one.

Our decomposition is a double decomposition because the U(1) four-vector field C_μ is also decomposed. This results to a Yang-Mills field with three dynamical degrees of freedom. In this decomposition, both monopoles and vortices appear. In addition, Dirac quantization condition is reproduced in the presence of vortices with a rescaled electric charge.

VI. ACKNOWLEDGMENTS

We are grateful to the research council of University of Tehran for supporting this study.

-
- [1] G. 't Hooft, Nucl. Phys. B **153**, 141 (1979); A. Polyakov, Nucl. Phys. B **120**, 429 (1977).
 - [2] Y. Nambu, Phys. Rev. D **10** 4262 (1974); M. Creutz, Phys. Rev. D **10** 2696 (1974); G. 't Hooft, High Energy Physics, Editorice Compositori, Bologna (1975); S. Mandelstam, Phys. Report **23** 245 (1976).
 - [3] G. Ripka, Dual Superconductor Models of Color Confinement, Springer-Verlag, 2005, arXiv: 0806.1078v2 [hep-th].
 - [4] G. 't Hooft, Nucl. Phys. B **190** 455 (1981).
 - [5] Y. M. Cho, Phys. Rev. D **21** (1980) 1080; D **23** 2415 (1981).
 - [6] L. Faddeev and A. J. Niemi, Phys. Rev. Lett. **82** 1624 (1999).

- [7] S. V. Shabanov, Phys. Lett. B **458** 322 (1999); Phys. Lett. B **463** 263 (1999).
- [8] T. Suzuki and I. Yotsuyanagi, Phys. Rev. D **42** 4257 (1990).
- [9] J. D. Stack, S. D. Neiman, and R. Wensley, Phys. Rev. D **50** 3399 (1994); H. Shiba and T. Suzuki, Phys. Lett. B **333** 461 (1994).
- [10] T. T. Wu and C. N. Yang, in Properties of Matter Under Unusual Conditions, edited by H. Mark and S. Fernbach (Interscience, New York, 1969).
- [11] T. Suzuki, Prog. Theor. Phys. **80** 929 (1988); H. Suganuma, S. Sasaki, and H. Toki, Nucl. Phys. B **435** 207 (1995).
- [12] L. S. Grigorio, M. S. Guimaraes, W. Oliveira, R. Rougemont, and C. Wotzasek, Phys. Lett. B **697** 392 (2011); S. Deldar and A. Mohamadnejad, Phys. Rev. D **86** 065005 (2012).
- [13] Y. M. Cho, H. W. Lee, and D. G. Pak, Phys. Lett. B **525** 347 (2002); W. S. Bae, Y. M. Cho, and S. W. Kimm, Phys. Rev. D **65** 025005 (2001).
- [14] L. Faddeev and A. J. Niemi, Phys. Lett. B **464** 90 (1999).
- [15] L. Faddeev and A. J. Niemi, Phys. Lett. B **525** 195 (2002).